LESSON MAKER PRO

Inverse Angle Lab

Use restricted trig domains to match each output with one input.

HIGH SCHOOL · HSF-TF.B.6

CCSS.Math.Content.HSF-TF.B.6 "(+) Understand that restricting a trigonometric function to a domain on which it is always increasing or always decreasing allows its inverse to be constructed."

NAME _____ DATE _____

ONE DIRECTION, ONE INVERSE

An inverse function needs each output to come from only one input. Restrict a trigonometric function to an interval where it moves in one direction only, always increasing or always decreasing. On that interval, you can reverse the input-output relationship to find inverse values.

WORKED EXAMPLE

Let f(x)=cos x with 0 <= x <= pi. Find $f^{-1}(-1/2)$.

1. On 0 <= x <= pi, cos x is always decreasing, so f has an inverse.
2. $f^{-1}(-1/2)$ is the x-value for which cos x=-1/2.
3. cos(2pi/3)=-1/2, and 2pi/3 is in the restricted domain.

Answer: $f^{-1}(-1/2)=2pi/3$

PRACTICE 6 problems

1 Let f(x)=sin x with -pi/2 <= x <= pi/2. Does f have an inverse? State the domain and range of f^{-1}.

ANSWER _____

2 Let g(x)=cos x with 0 <= x <= pi. Find $g^{-1}(-sqrt(3)/2)$.

ANSWER _____

3 Let h(x)=tan x with -pi/2 < x < pi/2. Find $h^{-1}(-sqrt(3))$.

ANSWER _____

4 Which restriction lets sin x have an inverse: [0, pi], [-pi/2, pi/2], or [pi/4, 5pi/4]? Explain briefly.

ANSWER _____

5 Let p(x)=cos x with -pi <= x <= 0. Find $p^{-1}(0)$.

ANSWER _____

6 Let q(x)=sin x with pi/2 <= x <= 3pi/2. Find $q^{-1}(-1)$.

ANSWER _____

APPLY IT Show your work

1 During one rising part of an audio visualizer, the screen height is modeled by h=sin t, where -pi/2 <= t <= pi/2. The screen shows h=-sqrt(2)/2. What is t? Why can the app use an inverse on this interval?

WORK SPACE

ANSWER _____

2 In a video game, a lighting effect is modeled by b=cos x, where 0 <= x <= pi. When b=-sqrt(2)/2, what is x? Explain why this restriction gives one angle for the brightness value.

WORK SPACE

ANSWER _____

CHALLENGE One step up

A student restricts r(x)=cos x to pi/6 <= x <= 11pi/6 and says r has an inverse because the interval is limited. Is the student correct? Use two different inputs from the interval to explain your reasoning.

WORK SPACE AND ANSWER

ANSWER KEY

Inverse Angle Lab

HIGH SCHOOL · HSF-TF.B.6

PART 1 · PRACTICE

#	PROBLEM	ANSWER
1	Let f(x)=sin x with -pi/2 <= x <= pi/2. Does f have an inverse? State the domain and range of f^{-1}.	**Yes. The domain of f^{-1} is [-1, 1], and the range of f^{-1} is [-pi/2, pi/2].**
2	Let g(x)=cos x with 0 <= x <= pi. Find g^{-1}(-sqrt(3)/2).	**g^{-1}(-sqrt(3)/2)=5pi/6**
3	Let h(x)=tan x with -pi/2 < x < pi/2. Find h^{-1}(-sqrt(3)).	**h^{-1}(-sqrt(3))=-pi/3**
4	Which restriction lets sin x have an inverse: [0, pi], [-pi/2, pi/2], or [pi/4, 5pi/4]? Explain briefly.	**[-pi/2, pi/2], because sin x is always increasing on this interval.**
5	Let p(x)=cos x with -pi <= x <= 0. Find p^{-1}(0).	**p^{-1}(0)=-pi/2**
6	Let q(x)=sin x with pi/2 <= x <= 3pi/2. Find q^{-1}(-1).	**q^{-1}(-1)=3pi/2**

PART 2 · APPLY IT

1. **t=-pi/4.** The app can use an inverse because sin t is always increasing on -pi/2 <= t <= pi/2. Solve sin t=-sqrt(2)/2 within the restricted interval. The restriction makes sine one-to-one because it is increasing throughout the interval.
2. **x=3pi/4.** Cosine is always decreasing on 0 <= x <= pi, so each brightness value comes from one angle. Solve cos x=-sqrt(2)/2 on the interval from 0 to pi. Cosine decreases on that interval, so the inverse is defined there.

CHALLENGE

No. cos(pi/3)=1/2 and cos(5pi/3)=1/2, but pi/3 and 5pi/3 are different inputs in the restricted interval. Since one output has two inputs, r is not one-to-one and has no inverse on this interval.

Accept equivalent interval notation and correct monotonicity explanations. Require exact radian values that lie in the stated restricted domain.