

Slope Match Mission

Find an instant that matches an interval's average rate of change.

HIGH SCHOOL • MA.912.C.2.10

MA.912.C.2.10 "Apply the Mean Value Theorem."

NAME _____

DATE _____

THE MEAN VALUE LINK

The Mean Value Theorem applies when f is continuous on $[a,b]$ and differentiable on (a,b) . It guarantees at least one number c in (a,b) where $f'(c) = (f(b)-f(a))/(b-a)$. First check the conditions, then find the average rate of change and solve for c .

WORKED EXAMPLE

Use the Mean Value Theorem to find c for $f(x)=x^2+13$ on $[5,11]$.

1. f is a polynomial, so it is continuous on $[5,11]$ and differentiable on $(5,11)$.
2. $(f(11)-f(5))/(11-5) = (134-38)/6 = 16$.
3. $f'(x)=2x$, so $2c=16$.
4. $c=8$, and 8 is in $(5,11)$.

Answer: $c=8$

PRACTICE 6 problems

- 1** For $f(x)=3x-2$ on $[0,4]$, find all values of c guaranteed by the Mean Value Theorem.

ANSWER _____

- 2** Use the Mean Value Theorem to find c for $f(x)=x^2-1$ on $[-2,2]$.

ANSWER _____

- 3** Use the Mean Value Theorem to find c for $f(x)=x^3$ on $[1,2]$.

ANSWER _____

- 4** Use the Mean Value Theorem to find c for $f(x)=1/x$ on $[1,4]$.

ANSWER _____

- 5** Does the Mean Value Theorem apply to $f(x)=|x|$ on $[-3,1]$? Explain briefly.

ANSWER _____

- 6** Use the Mean Value Theorem to find c for $f(x)=\sin x$ on $[0,\pi]$.

ANSWER _____

APPLY IT Show your work

- 1** During a 4-hour bike ride, Jordan's distance from the start, in miles, is modeled by $s(t)=t^2+2t$ for $0 \leq t \leq 4$. Use the Mean Value Theorem to find a time when Jordan's instantaneous velocity equals the average velocity for the entire ride.

WORK SPACE

ANSWER _____

- 2** A game download has $q(t)=t^2+1$ gigabytes downloaded t hours after it starts, for $1 \leq t \leq 3$. Find a time when the instantaneous download rate equals the average download rate over that interval.

WORK SPACE

ANSWER _____

CHALLENGE One step up

CHALLENGE

For $f(x)=x^3-3x$ on $[-2,2]$, use the Mean Value Theorem to find all values of c . Explain why each value works.

PART 1 · PRACTICE

#	PROBLEM	ANSWER
1	For $f(x)=3x-2$ on $[0,4]$, find all values of c guaranteed by the Mean Value Theorem.	Every c with $0 < c < 4$.
2	Use the Mean Value Theorem to find c for $f(x)=x^2-1$ on $[-2,2]$.	$c=0$
3	Use the Mean Value Theorem to find c for $f(x)=x^3$ on $[1,2]$.	$c=\sqrt[3]{7/3}$
4	Use the Mean Value Theorem to find c for $f(x)=1/x$ on $[1,4]$.	$c=2$
5	Does the Mean Value Theorem apply to $f(x)= x $ on $[-3,1]$? Explain briefly.	No. f is not differentiable at $x=0$, which is inside $(-3,1)$.
6	Use the Mean Value Theorem to find c for $f(x)=\sin x$ on $[0,\pi]$.	$c=\pi/2$

PART 2 · APPLY IT

- $t=2$ hours** The average velocity is $(s(4)-s(0))/(4-0)=6$ miles per hour. Since $s'(t)=2t+2$, solve $2t+2=6$ to get $t=2$.
- $t=2$ hours** The average download rate is $(q(3)-q(1))/(3-1)=4$ gigabytes per hour. Since $q'(t)=2t$, solve $2t=4$ to get $t=2$.

CHALLENGE

f is a polynomial, so the theorem applies. The average rate of change is $(f(2)-f(-2))/(2-(-2))=(2-(-2))/4=1$. Since $f'(x)=3x^2-3$, solve $3c^2-3=1$ to get $c=\pm\sqrt{4/3}$. Both values are in $(-2,2)$.

Accept $\sqrt{7/3}$ or $\sqrt{21}/3$. Accept $\pm\sqrt{2/3}$ or $\pm\sqrt{6}/3$ for the challenge, and accept any equivalent statement that every interior point works for the linear function.